## Problem Set 5

## Discrete Mathematics

Due on the 27<sup>th</sup> of February, 2024

(10 pts) 1. Find and explain the flaw(s) in this argument.

> We prove every nonempty set of people all have the same age. **Proof.** We denote the age of a person p by  $\alpha(p)$ .

Basis Step:

Suppose  $P = \{p\}$  is a set with one person in it. Clearly, all the people in *P* have the same age as each other.

Inductive Step:

Let  $k \in \mathbb{N}_+$  and suppose any set of k-many people all have the same age. Let  $P = \{p_1, p_2, \dots p_k, p_{k+1}\}$  be a set with k+1 people in it. Consider  $L := \{p_1, \dots p_k\}$  and  $R := \{p_2, \dots p_{k+1}\}$ . Since Land R both have k people, we know everyone in these sets has the same age by the inductive hypothesis.

Let  $\ell, r \in P$ . If  $\ell \in L \land r \in L$ , then  $\alpha(\ell) = \alpha(r)$ . Similarly, if  $\ell \in R \land r \in R$ , then  $\alpha(\ell) = \alpha(r)$ . Now, suppose  $\ell \in L \land r \in R$ .

$$\alpha(\ell) = \alpha(p_1) = \alpha(p_2) = \alpha(p_{k+1}) = \alpha(r)$$

So, all people in P have the same age.

Therefore, everyone on Earth has the same age. Q.E.D.

- 2. Show that  $\forall x (x \neq x \cup \{x\})$ . (20 pts)
- 3. We will work up to a proof of the commutativity of addition on  $\mathbb{N}$ . (15 pts)
  - (a) Show  $(\forall x \in \mathbb{N})(x+0=0+x)$ .
  - (b) Show  $(\forall x, y \in \mathbb{N})(x + \mathfrak{s}(y) = \mathfrak{s}(y) + x)$ .
  - (c) Show  $(\forall x, y \in \mathbb{N})(x + y = y + x)$ .
- 4. Show  $(\forall x, y, z \in \mathbb{N})(x \cdot (y+z) = (x \cdot y) + (x \cdot z))$ . (15 pts)
- 5. For this problem, you may assume the commutativity and associa-(20 pts) tivity of addition and multiplication over N. You may also assume multiplication distributes over addition on N. Prove the following statement for all  $n \in \mathbb{N}$ .

$$1 + \sum_{i=0}^{n} 2^{i} = 2^{n+1}$$

6. We say x is  $\in$ -transitive by definition when  $(\forall y \in x)(\forall z \in y)(z \in x)$ . (20 pts) Show that every natural number is  $\in$ -transitive.

Recall that the natural numbers are defined recursively as follows.

$$0 := \emptyset$$

$$\mathfrak{s}(n) := n \cup \{n\}$$

Addition on N is defined below.

$$n + 0 := n$$
  
 $n + \mathfrak{s}(m) := \mathfrak{s}(n + m)$ 

Multiplication on  $\mathbb N$  is defined below.

$$n \cdot 0 := 0$$
$$n \cdot \mathfrak{s}(m) := (n \cdot m) + n$$

Exponentiation on  $\mathbb N$  is defined below.

$$n^0 := 1$$

$$n^{\mathfrak{s}(m)} := n \cdot n^m$$

We define the iterated sum of a sequence of terms f(0), f(1), f(2), ... as follows.

$$\sum_{i=0}^{0} f(i) := f(0)$$

$$\sum_{i=0}^{\mathfrak{s}(n)} f(i) := \left(\sum_{i=0}^{n} f(i)\right) + f(\mathfrak{s}(n))$$

You may rely on the following theorems:  $(\forall x \in \mathbb{N})(\mathfrak{s}(x) = x + 1).$ 

$$(\forall x \in \mathbb{N})(\mathfrak{g}(x) = x + 1)$$

$$(\forall x \in \mathbb{N})(\mathfrak{s}(x) = 1 + x).$$

$$(\forall x, y, z \in \mathbb{N})(x + (y + z) = (x + y) + z).$$